

SPECIFICATION

 $\gamma\delta$ T CELL IMMUNOACTIVITY ENHANCERS CONTAINING
EXTRACT OF LENTINUS EDODES MYCELIIUM

Insa!

FIELD OF THE INVENTION

5 The present invention relates to the development and preparation of $\gamma\delta$ T cell activity enhancers and therefore immunopotentiators containing an extract of Lentinus edodes mycelium.

10 The present invention also relates to the development and preparation of foods, drinks and feeds containing an extract of Lentinus edodes mycelium and having a $\gamma\delta$ T cell activity-enhancing effect and therefore an immunopotentiating effect.

15 The present invention also relates to the development and preparation of antitumor agents, therapeutic agents against bacterial infections and therapeutic agents against viral infections containing an extract of Lentinus edodes mycelium.

20 PRIOR ARTCharacteristics of $\gamma\delta$ T cells

Peripheral T cells in animal blood are mainly classified into two groups based on the type of their cell surface antigens referred to as T cell receptors (TCR).

25 One type is an $\alpha\beta$ T cell bearing TCR α and β chains on their cell surfaces, and the other is a $\gamma\delta$ T cell bearing TCR γ and δ chains. $\gamma\delta$ T cells are cytotoxic killer cells present at a level of only a few to about 10% in normal peripheral blood

and lymphoid tissue in humans, and the characteristics of which are quite distinct from those of $\alpha\beta$ T cells.

In humans, $\gamma\delta$ T cells are present in the intestinal tract, skin and peripheral blood or the like and elicit a local immunity. So far, $\gamma\delta$ T cells have been reported to have functions such as cytotoxicity against cancer cells, and protective activity against bacterial or virus infections, etc.

10 Phylaxis activity of $\gamma\delta$ T cells

Some $\gamma\delta$ T cells in the spleen and other organs produce cytokines such as IL-4 or IFN- α in response to infection. It has been shown under experimental conditions that when these cells are lacking, resistance to bacterial infections is reduced. For example, there is one report which describes that resistance to Mycobacterium tuberculosis infection decreased in mice treated with a $\gamma\delta$ -type TCR antibody to transiently inhibit functions of $\gamma\delta$ T cells or mice deficient in the TCR γ gene (Ladel C. et al., Eur. J. Immunol., 1995, 25:2877-2881). Another report describes that $\gamma\delta$ T cells appear during the early stage of infection with Listeria monocytogenes (Hiromatsu K. et al., J. Exp. Med., 1992, 175:49-56). These findings suggest that $\gamma\delta$ T cells play an important role in protecting against bacterial infections.

It has also been reported that chronic hepatitis B virus infection induces the growth of $\gamma\delta$ T cells in the liver and spleen (Ozaki S. et al., J. Med. Invest., 1998,

44:215-217); and that vaccinia virus markedly increases during the early stage of infection in $\gamma\delta$ T cell-deficient mice as compared with normal mice (Welsh RM, et al., Immunol. Rev., 1997, 159:79-93). These findings suggest
5 that $\gamma\delta$ T cells act not only on bacterial infections but also on viral infections.

Cytotoxicity of $\gamma\delta$ T cells against cancer cells

$\gamma\delta$ T cells are a class of T cells that are capable of
10 specifically targeting and killing autologous cancer cells, but show no cytotoxicity to autologous normal lymphocytes (such as $\alpha\beta$ T cells). In this respect, in cancer therapy which employs activated $\gamma\delta$ T cells there is very little danger of side effects. In contrast to $\gamma\delta$ T cells, $\alpha\beta$ T
15 cells are known to kill autologous leukocytes rather than autologous cancer cells, and thus in cancer therapy which employs activated $\alpha\beta$ T cells there is a likelihood that serious side effects will occur. In view of this, cancer therapy which employs activated $\gamma\delta$ T cells is desirable.

20 Moreover, $\gamma\delta$ T cells have similar characteristics to NK cells such as their MHC-nonrestricted cytotoxicity against cancer cells. $\gamma\delta$ T cells are present in peripheral blood of children at about 10% but decrease with age. This suggests that the increase in the occurrence of cancer with
25 age may be related to a decrease in $\gamma\delta$ T cells. In peripheral blood of chicken, sheep, cow or the like, $\gamma\delta$ T cells are found at levels as high as 15-50%. The low incidence of tumors in these animals suggests that the

presence of peripheral $\gamma\delta$ T cells may contribute greatly to the inhibition of cancer.

Pharmacological effects of Lentinus edodes

Lentinus edodes (Shiitake) is a common edible mushroom found in both Japan and China, and has been cultivated in Japan for around 300 years. The part of the mushroom used as food consists of the reproductive body, also referred to as the fruiting body of fungi, which produces spores for reproduction, while the vegetative body includes hyphae which produce mycelia extending into a growing area such as soil or logs.

Lentinus edodes has long been believed to have some effect against a variety of diseases and symptoms, but it is only relatively recently that any pharmacological effect has been elucidated. Various effects of extract of Lentinus edodes mycelium are reported, these include the inhibition of oncogenesis in the large bowel and liver and the growth of transplanted tumor cells and increased survival of animals in carcinogenesis experiments in rats and mice (N. Sugano et al., Cancer Letter, 27:1, 1985; Y. Suzuki et al., Journal of the Japan Society of Coloproctology, 43:178, 1990, etc.); mitogenic activity (T. Tabata et al., Immunopharmacology, 24:57, 1992; Y. Hibino et al., Immunopharmacology, 28:77, 1994, etc.); enhanced antibody production and inhibitory effects against immunological hepatocyte damage caused by ADCC (antibody-dependent cell-mediated cytotoxicity) (Y. Mizoguchi et al., Journal of Hepato-Biliary-Pancreatic Study, 15:127, 1987).

09886717.021902

These findings have acted as a catalyst for concentrated research into the pharmacological effects of ingredients of *Lentinus edodes* in the medical and pharmaceutical field. As a result, researchers have
5 discovered that some of the constituents of *Lentinus edodes* can be used in the treatment of cancer or other diseases by restoring immune function in humans, and further that such constituents may also inhibit the onset of cancer.

In accomplishing the present invention the inventors
10 have aimed to further elucidate pharmacological effects of an extract of *Lentinus edodes* mycelium, and also to search for new applications of the extract in the form of drugs, foods, drinks, feeds, etc.

One object of the present invention is to develop and
15 provide a $\gamma\delta$ T cell activity enhancer and therefore an immunopotentiator such as an antitumor agent, therapeutic agent against bacterial infections and therapeutic agent against viral infections containing an extract of *Lentinus edodes* mycelium.

20 Another object of the present invention is to use a $\gamma\delta$ T cell activity enhancer containing an extract of *Lentinus edodes* mycelium or an immunopotentiator containing an extract of *Lentinus edodes* mycelium to treat a tumor in a subject.

25

DISCLOSURE OF THE INVENTION

As a result of intensive studies aimed at solving the above problems, the present invention has been accomplished

on the basis of the finding that an extract of Lentinus edodes mycelium exhibits an effect of remarkably enhancing the activity of $\gamma\delta$ T cells.

Accordingly, the present invention has been developed and provided $\gamma\delta$ T cell activity enhancers and therefore immunopotentiators such as antitumor agents or therapeutic agents against bacterial or viral infections containing the extract of Lentinus edodes mycelium.

The present invention also provides methods developed for treating tumor, bacterial infections and viral infections using the extract of Lentinus edodes mycelium.

$\gamma\delta$ T cell activity enhancers or immunopotentiators of the present invention may be in the form of a pharmaceutical composition containing the extract of Lentinus edodes mycelium and optionally a pharmaceutically acceptable carrier.

$\gamma\delta$ T cell activity enhancers or immunopotentiators of the present invention may be administered in the form of injection or oral, mucosal, gastrointestinal or percutaneous formulation.

$\gamma\delta$ T cell activity enhancers or immunopotentiators of the present invention may also be in the form of a food, drink or feed.

25 BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 shows that the proportion of $\gamma\delta$ T cells in peripheral blood increases following administration of the extract of Lentinus edodes mycelium of the present

invention.

FIG. 2 shows the results of flow cytometric analysis of $\gamma\delta$ T cells before and after administration of the extract of *Lentinus edodes* mycelium.

5 FIG. 3 shows that the proportion of $\alpha\beta$ T cells in peripheral blood decreases following administration of the extract of *Lentinus edodes* mycelium of the present invention.

10 THE MOST PREFERRED EMBODIMENTS OF THE INVENTION

206430-17-04130
An extract of *Lentinus edodes* mycelium used for enhancing the activity of $\gamma\delta$ T cells according to the present invention refers to an extract obtained by crushing and decomposing mycelia grown from *Lentinus edodes* cultured
15 on a solid medium, or a solid medium itself containing *Lentinus edodes* mycelia in the presence of water and an enzyme.

An extract of *Lentinus edodes* mycelium used herein is preferably obtained by, but not limited to, the following
20 process. *Lentinus edodes* spawn is inoculated on a solid medium based on bagasse (sugar cane residue) and defatted rice bran to grow mycelia, and then the solid medium containing the grown mycelia is delignified so that 30% by weight or less is able to pass through a 12-mesh sieve. To
25 this delignified solid medium are added water and one or more enzymes selected from cellulase, protease or glucosidase while maintaining said solid medium at a temperature of 30-55°C, and said solid medium is crushed

and ground in the presence of said enzyme so that at least 70% by weight of bagasse fiber is able to pass through a 12-mesh sieve. Then, the temperature is raised to 95°C to ensure inactivation of the enzyme and sterilization, and the resulting suspension is filtered to give an extract of *Lentinus edodes* mycelium. The extract of *Lentinus edodes* mycelium as prepared above may be directly used in γ DT cell activity enhancers of the present invention, but conveniently concentrated and freeze-dried into powder to be stored and used in various forms. The freeze-dried product is a brown powder with hygroscopic characteristics and has a peculiar taste and odor.

The extract of *Lentinus edodes* mycelium was tested for its in vivo $\gamma\delta T$ cell activity-enhancing effect by the method described in the examples below, and it will be seen from these examples that it has a remarkable in vivo $\gamma\delta T$ cell activity-enhancing effect.

20 $\gamma\delta$ T cell activity enhancers of the present invention are effective for treating and/or preventing tumor induced by tumor cells to which $\gamma\delta$ T cells are cytotoxic. $\gamma\delta$ T cell activity enhancers of the present invention are characterized in that they enhance the activity of $\gamma\delta$ T cells and thereby lead to the destruction of tumor cells under the action of the activated $\gamma\delta$ T cells, rather than
25 having a direct action on specific tumor cells. Consequently, tumor cells to be treated with $\gamma\delta$ T cell activity enhancers of the present invention may be not only malignant tumor cells but also benign tumor cells, and are

not limited to specific tumor cells. In addition, $\gamma\delta$ T cell activity enhancers containing the extract of *Lentinus edodes* mycelium and optionally a pharmaceutically acceptable carrier can be used in the form of both therapeutic and prophylactic compositions.

10 $\gamma\delta$ T cell activity enhancers of the present invention
 can also be used as therapeutic and/or prophylactic
 compositions for bacterial or viral infections. $\gamma\delta$ T cell
 activity enhancers of the present invention are intended to
 enhance the activity of $\gamma\delta$ T cells and thereby to remove
 infecting bacteria or viruses from the patient rather than
 to directly act on specific bacteria or viruses. Bacterial
 or viral diseases that can be treated with $\gamma\delta$ T cell
 activity enhancers of the present invention include such
 15 diseases as , but are not limited to, those caused by
 Mycobacterium spp. Listeria monocytogenes, hepatitis
 viruses (types A, B and C), human immunodeficiency virus,
 vaccinia virus and the like.

γδT cell activity enhancers of the present invention
in the form of a therapeutic and/or prophylactic
composition are administered most preferably via the oral
route, but may also be administered via intravenous,
intraperitoneal, subcutaneous, intramuscular, nasal,
percutaneous or other route. Dosage forms suitable for
oral administration include, but are not limited to,
tablets, capsules, powders, granules, solutions, syrups,
etc. Dosage forms suitable for nasal or percutaneous
administration include, but not limited to, cataplasms,

p

patches, etc.

Pharmaceutically acceptable carriers include, but are not limited to, suitable excipients, binders, disintegrators, lubricants, flavoring agents, colorants, solubilizers, suspending agents, coating agents or the like known in the art.

Pharmaceutically acceptable carriers that can be optionally mixed in $\gamma\delta$ T cell activity-enhancing formulations of the present invention include, but are not limited to excipients such as lactose, dextrose, starch, crystalline cellulose; binders such as starch, gelatin, methyl cellulose, polyvinylpyrrolidone; disintegrators such as starch, calcium carboxymethylcellulose, carboxymethyl starch; lubricants such as talc, stearates; coating agents such as sucrose, talc, gelatin; as well as various brighteners, flavoring agents, colorants, corrigents, solubilizers, stabilizers, suspending agents, absorbefaciants or the like known in the art depending on the purpose. For use as injections, various diluents commonly used in this field of art such as water or ethyl alcohol can be used.

The dose of $\gamma\delta$ T cell activity enhancers of the present invention is determined by physicians taking into account the age, weight and condition of the subject, the route of administration and other factors. The dose is not strictly limited because the extract of *Lentinus edodes* mycelium contained in $\gamma\delta$ T cell activity enhancers of the present invention is highly safe, and has been

traditionally ingested as a component of food. For example, the extract of *Lentinus edodes* mycelium is normally administered preferably at a dose of 100 mg - 10000 mg several times (about 2-3 times) daily (a total of 200 mg - 30000 mg daily), more preferably 500 mg - 5000 mg three times daily (a total of 1500 mg - 15000 mg daily), most preferably 1000 mg - 1500 mg three times daily (a total of 3000 mg - 4500 mg daily). It may be administered in combination with other antitumor agents.

5
10 $\gamma\delta$ T cell activity enhancers of the present invention can be provided in a dosage form also suitable for adoptive immunotherapy for treating tumor. Adoptive immunotherapy refers to a type of anti-tumor therapy intended to kill tumor cells by transferring into a subject sensitized cells, normally lymphocytes. In the case of the present invention, $\gamma\delta$ T cells are initially isolated from peripheral blood of the subject, and the isolated $\gamma\delta$ T cells are activated in vitro by a $\gamma\delta$ T cell activity enhancer of the present invention, and then the activated $\gamma\delta$ T cells are returned into the subject. As a result, tumor cells in the subject can be destroyed by the action of the activated $\gamma\delta$ T cells.

15 $\gamma\delta$ T cell activity enhancers of the present invention may be the extract of *Lentinus edodes* mycelium itself or pharmaceutical or veterinary compositions comprising a $\gamma\delta$ T cell activity enhancer containing the extract of *Lentinus edodes* mycelium and a pharmaceutically acceptable carrier.

25 $\gamma\delta$ T cell activity enhancers of the present invention can also be provided in the form of a food. Preferred

09856717 "021902

forms of food include powders, granules, pastes, jellies, etc. Granules desirably are supplemented with sugars such as lactose to add a sweet taste. $\gamma\delta$ T cell activity enhancers of the present invention can also be provided in the form of a drink. These foods or drinks may be supplemented with vitamins, minerals such as calcium, alcohols, deodorants such as polyphenols in addition to the extract of *Lentinus edodes* mycelium. These foods or drinks include the categories of specific health foods, medical foods or the like.

$\gamma\delta$ T cell activity enhancers of the present invention can also be provided in the form of a feed or feed additive. $\gamma\delta$ T cell activity enhancers of the present invention can be used as a feed or feed additive for domestic animals to treat and/or prevent tumor occurring in domestic animals or to treat and/or prevent bacterial or viral infections in domestic animals. As a result, the amount of therapeutic agents such as antibiotics currently used can be reduced, thereby reducing farming costs. Another advantage is that the period during which shipment of animals is suspended due to the administration of antibiotics can be shortened.

In vivo $\gamma\delta$ T cell activity-enhancing effect was tested in human subjects as follows. Human subjects initially received 3.6 g of the extract of *Lentinus edodes* mycelium bulk powder daily for 7 days (a total of 25.2 g). Then, the proportion of $\gamma\delta$ T cells in peripheral blood after administration of the extract of *Lentinus edodes* mycelium was determined by flow cytometry as compared with the

proportion before administration.

The following examples, which further illustrate the present invention, should not be taken as limiting the the scope of the invention thereto. Various changes and
5 modifications can be made by those skilled in the art and such changes and modifications are also included in the scope of the present invention.

EXAMPLES

10 Example 1: Preparation of an extract of *Lentinus edodes* mycelium

A solid medium consisting of 90 parts by weight of bagasse and 10 parts by weight of rice bran was soaked with an appropriate amount of pure water, and then inoculated
15 with *Lentinus edodes* spawn and allowed to stand in an incubator at controlled temperature and humidity to grow mycelia. After mycelia spread over the solid medium, the bagasse base was delignified so that 24% by weight or less may pass through a 12-mesh sieve. To 1.0 kg of this
20 delignified medium were added 3.5 L of pure water and 2.0 g of purified cellulase while maintaining the solid medium at 40°C to prepare a medium-containing mixture.

Then, the medium-containing mixture was circulated by a variable speed gear pump, during which the solid medium
25 was crushed and ground under the gears for about 200 minutes so that about 80% by weight of bagasse fiber may pass through a 12-mesh sieve. The medium-containing mixture was crushed and ground while the temperature of

said mixture was gradually increased. Then, the medium-containing mixture was further heated to 90°C and allowed to stand for 30 minutes. This heating to 90°C ensured inactivation of the enzyme and sterilization. The

5 resulting medium-containing mixture was filtered through a 60-mesh filter cloth to give an extract of *Lentinus edodes* mycelium solution, which was concentrated and then freeze-dried into an extract of *Lentinus edodes* mycelium bulk powder.

10 The extract of *Lentinus edodes* mycelium as prepared above contained 25.3% (w/w) carbohydrates determined by the phenol-sulfuric acid method, 19.7% (w/w) proteins determined by the Lowry method and 2.6% (w/w) polyphenols determined by the Folin-Denis method using gallic acid as
15 standard. The extract of *Lentinus edodes* mycelium further contains 8% crude fat, 22% crude ash and about 20% soluble nitrogen-free materials other than carbohydrates.

The extract of *Lentinus edodes* mycelium had a sugar composition (%) as follows: Xyl 15.2, Ara 8.2, Man 8.4, Gul
20 39.4, Gal 5.4, GlcN 12.0, GluUA 11.3.

Example 2: In vivo $\gamma\delta$ T cell activity-enhancing test of the extract of *Lentinus edodes* mycelium

Three human subjects (subjects A-C) received orally
25 3.6 g/day of the extract of *Lentinus edodes* mycelium bulk powder daily for 7 days (a total of 25.2 g). After the period of administration of the extract of *Lentinus edodes* mycelium, peripheral blood was collected from the human

subjects. The proportion of $\gamma\delta$ T cells in peripheral blood collected after administration was determined by flow cytometry as compared with the proportion in peripheral blood collected before administration. The results are shown in Figs. 1 and 2.

In all of the three subjects, the proportion of $\gamma\delta$ T cells in peripheral blood increased by an average of 40% or more after administration of the extract of *Lentinus edodes* mycelium as compared with before administration.

10

Table 1: Increase of the proportion of $\gamma\delta$ T cells in peripheral blood after administration

	Subject A	Subject B	Subject C	Mean \pm SEM
Increase	124.39%	146.15%	150.00%	140.18% \pm 7.97%

15

Peripheral blood collected from the same subject before and after administration was tested for other markers than $\gamma\delta$ T cells. As a result, the extract of *Lentinus edodes* mycelium showed no proliferative activity on $\alpha\beta$ T cells and on average the proportion of $\alpha\beta$ T cells decreased (Fig. 3).

20

INDUSTRIAL APPLICABILITY

$\gamma\delta$ T cell activity enhancers containing the extract of *Lentinus edodes* mycelium of the present invention were found to actually activate $\gamma\delta$ T cells. Thus, $\gamma\delta$ T cell activity enhancers of the present invention can be used for preventing or treating tumor, bacterial infections and

25

09856717 "021902

viral infections because they have the effect of protecting living bodies against tumor, bacterial infections and viral infections by inducing cytotoxic activity against tumor cells, antibacterial activity and antiviral activity of $\gamma\delta$ T cells. Moreover, $\gamma\delta$ T cell activity enhancers of the present invention are suitable for wide industrial application because they can be used safely without side effects.

They can also be used in domestic animals with bacterial and/or viral infections to reduce the amount of therapeutic agents currently used such as antibiotics, thereby reducing costs for raising. They also have the advantage that the period during which shipment is suspended can be shortened because antibiotics are not used.